

PEDIATRIC NEUROLOGY BRIEFS

A MONTHLY JOURNAL REVIEW

J. GORDON MILLICHAP, M.D., F.R.C.P., EDITOR

Vol. 24, No. 8

August 2010

VASCULAR DISORDERS

SURGICAL OUTCOME OF MOYAMOYA DISEASE

Researchers at Seoul National University Children's Hospital, Korea, analyzed the clinical characteristics and prognostic factors for outcome of 410 pediatric cases of moyamoya disease (MMD) after bypass surgery. Patients' ages were 0-3 years in 15%, 4-6 in 27%, and 7-18 years in 58%; sex M/F ratio was 1:1.3. Cases were familial in 12%. Clinical manifestations included transient ischemic attacks in 56%, infarction in 39%, headache in 33%, seizure 19%, involuntary movements 4%, and hemorrhage 3%. Unilateral MMD at initial diagnosis in 53 progressed to bilateral involvement in 24 (45%) within 2 years. CT/MRI showed no infarction in 44%, and major infarction in 39%. SPECT showed decreased perfusion in 60% and decreased vascular reserve in 84%. Neurocognition test was abnormal in 93%, and IQ <90 in 24% (>90 in 76%). Surgery consisted of bilateral encephaloduroarteriosynangiosis augmented by bifrontal encephalogaleo/periosteal synangiosis. The interval between onset of symptoms and surgery averaged 21 months (range, 1 week to 108 months). In cases benefited, symptoms usually resolved in an average of 4.3 months. Surgical outcome was excellent in 66%, good in 15%, fair in 15%, and poor in 4% of patients. Infarction at presentation was associated with unfavorable clinical outcome, and decreased vascular reserve only on SPECT indicative of a favorable outcome (81% cases). Early diagnosis and surgical intervention before irreversible hemodynamic change occurs are essential to obtain a favorable outcome. (Kim S-K, Cho B-K, Phi JH, et al. Pediatric moyamoya disease: an analysis of 410 consecutive cases. *Ann Neurol* July 2010;68:92-101). (Respond: Dr K-C Wang, Division of Pediatric Neurosurgery, Seoul National University Children's Hospital, Seoul 110-744, Republic of Korea. E-mail: kewang@snu.ac.kr).

PEDIATRIC NEUROLOGY BRIEFS (ISSN 1043-3155) © 2010 covers selected articles from the world literature and is published monthly. Send subscription requests (\$68 US; \$72 Canada; \$75 airmail outside N America) to **Pediatric Neurology Briefs - J. Gordon Millichap, M.D., F.R.C.P.-Editor**, P.O. Box 11391, Chicago, Illinois, 60611, USA. The editor is Pediatric Neurologist at Children's Memorial Hospital and Professor Emeritus, Northwestern University Medical School, Chicago, Illinois.

PNB is a continuing education service designed to expedite and facilitate review of current scientific information for physicians and other health professionals. Fax: 312-943-0123.

COMMENT. Moyamoya disease is a largely Korean or Japanese cerebrovascular disease of internal carotid arteries with compensatory vascular network at the base of the brain. Repeated transient ischemic attacks are common presenting symptoms in children and intracranial hemorrhage in adults. Surgery to establish adequate collateral circulation is effective in ischemic cases without infarction. Outcome was favorable in 81% of the above large series of pediatric cases. Infarction at the time of diagnosis of MMD is predictive of a 3-fold increased risk of unfavorable surgical outcome, whereas decreased vascular reserve only with normal basal perfusion on SPECT has a 14-fold increased frequency of favorable outcome. Long-term follow-up is recommended in unilateral cases to monitor potential progression and bilateral involvement.

DIAGNOSIS AND TREATMENT OF VEIN OF GALEN ANEURYSMAL MALFORMATIONS

Diagnosis and treatment of 13 patients with vein of Galen aneurysmal malformation (VGAM), seen at Children's Hospital and University of Pennsylvania, Philadelphia, 1994-2007, were evaluated retrospectively, and those treated within the first 2 weeks were compared with those treated later. Six (46%) patients were diagnosed in utero by fetal ultrasound or fetal MR, three within 1 week of birth with cardiac failure, and four from 1.5 to 31 months of age. Cardiac failure was the most common presenting symptom, diagnosed in the first week of birth, and was present in 7 patients. Increasing head circumference, hydrocephalus and developmental delay characterized all but one of the patients presenting outside the neonatal period. One patient presented with subarachnoid hemorrhage at 31 months of age. Postnatal MRAs demonstrated preembolization hemorrhage in 4 patients, ischemia in 4, and diffuse parenchymal abnormality in 5. Five had hydrocephalus. Angioarchitecture of a choroidal pattern was demonstrated in 62% of VGAMs, and in all VGAMs presenting early with cardiac failure. Five of 6 presenting later in life had mural angioarchitecture. Endovascular treatment was implemented in 11 patients. Of 6 patients treated emergently in the first 2 weeks, 2 developed normally, 1 had mild to moderate neurological deficits, 1 had severe deficits, and 2 died. Outcome in 5 patients treated later, at 1.5 to 31 months, was better, normal in 3 and mild neurological deficits in 2. (Heuer GG, Gabel B, Beslow LA, et al. Diagnosis and treatment of vein of Galen aneurysmal malformations. *Childs Nerv Syst* July 2010;26:879-887). (Respond: Dr RW Hurst, Division of Neuroradiology, Hospital of the University of Pennsylvania, 3400 Spruce Street, Philadelphia, PA 19104. E-mail: Robert.Hurst@uphs.upenn.edu).

COMMENT. Cardiac failure is the presenting feature of neonatal cases of VGAM, and enlarging head circumference, hydrocephalus and developmental delay characterize cases with later onset. Endovascular management is the preferred approach in all age groups. Outcome in cases presenting after the neonatal period is better than those of early onset. Cardiac failure is particularly intractable, with mortality rates as high as 62%. In utero MR diagnosis allows for early stabilization and cardiac management.